

FILE 'HCAPLUS' ENTERED AT 16:30:39 ON 25 AUG 2009

L1 17172 S ANTIPSYCHOTIC OR RISPERIDONE OR OLANZAPINE OR QUETIAPINE OR Z
L2 32609 S ANTIDEPRESSANT OR (SELECTIVE SEROTONIN REUPTAKE INHIBITOR) OR
L3 6612 S ESCITALOPRAM OR BUPROPION OR NEFAZODONE OR MIRTAZAPINE OR VEN
L4 8608 S SUICIDE OR SUICIDAL OR SUICIDALITY
L5 612 S L1 AND L2 AND (L3 OR L4)
L6 148 S L5 AND (PY<2003 OR AY<2003 OR PRY<2003)

FILE 'STNGUIDE' ENTERED AT 16:31:35 ON 25 AUG 2009

FILE 'HCAPLUS' ENTERED AT 16:32:06 ON 25 AUG 2009

L7 99340 S DEPRESSION OR DEPRESSIVE
L8 51 S L6 AND L7

=> file hcaplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	1.76	1.76

FILE 'HCAPLUS' ENTERED AT 16:30:39 ON 25 AUG 2009
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 25 Aug 2009 VOL 151 ISS 9
 FILE LAST UPDATED: 24 Aug 2009 (20090824/ED)
 REVISED CLASS FIELDS (/NCL) LAST RELOADED: Jun 2009
 USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Jun 2009

HCAplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

The ALL, BIB, MAX, and STD display formats in the CA/CAPLUS family of databases have been updated to include new citing references information. This enhancement may impact record import into database management software. For additional information, refer to NEWS 9.

=> s antipsychotic or risperidone or olanzapine or quetiapine or ziprasidone or aripiprazole or iloperidone or melperone or amperozide or perphenazine or trifluoroperazine or zotepine or flupenthixol or amisulpride or sulpride

12384 ANTIPSYCHOTIC
 3602 RISPERIDONE
 3250 OLANZAPINE
 1653 QUETIAPINE
 1084 ZIPRASIDONE
 1038 ARIPIPAZOLE
 110 ILOPERIDONE
 195 MELPERONE
 157 AMPEROZIDE
 1771 PERPHENAZINE
 481 TRIFLUOROPERAZINE
 291 ZOTEPINE
 6 FLUPENTHIXOL
 478 AMISULPRIDE
 27 SULPRIDE

L1 17172 ANTIPSYCHOTIC OR RISPERIDONE OR OLANZAPINE OR QUETIAPINE OR ZIPRASIDONE OR ARIPIPRAZOLE OR ILOPERIDONE OR MELPERONE OR AMPEROZIDE OR PERPHENAZINE OR TRIFLUOROPERAZINE OR ZOTEPINE OR FLUPHENTHIXOL OR AMISULPRIDE OR SULPRIDE

=> s antidepressant or (selective serotonin reuptake inhibitor) or SSRI or fluoxetine or norfluoxetine or paroxetine or sertaline or fluvoxamine or citalopram

24648 ANTIDEPRESSANT
493624 SELECTIVE
78418 SEROTONIN
12126 REUPTAKE
624610 INHIBITOR
2079 SELECTIVE SEROTONIN REUPTAKE INHIBITOR
(SELECTIVE(W)SEROTONIN(W)REUPTAKE(W)INHIBITOR)
2153 SSRI
7121 FLUOXETINE
501 NORFLUOXETINE
3971 PAROXETINE
8 SERTALINE
2242 FLUVOXAMINE
3396 CITALOPRAM

L2 32609 ANTIDEPRESSANT OR (SELECTIVE SEROTONIN REUPTAKE INHIBITOR) OR SSRI OR FLUOXETINE OR NORFLUOXETINE OR PAROXETINE OR SERTALINE OR FLUVOXAMINE OR CITALOPRAM

=> s escitalopram or bupropion or nefazodone or mirtazapine or venlafaxine or duloxetine or milnacipran or reboxetine zimelidine or indalpine or gepirone or femoxetine or alaproclate

723 ESCITALOPRAM
1896 BUPROPION
821 NEFAZODONE
1059 MIRTAZAPINE
2267 VENLAFAXINE
947 DULOXETINE
513 MILNACIPRAN
672 REBOXETINE
452 ZIMELIDINE
0 REBOXETINE ZIMELIDINE
(REBOXETINE(W)ZIMELIDINE)
142 INDALPINE
359 GEPIRONE
152 FEMOXETINE
171 ALAPROCLATE

L3 6612 ESCITALOPRAM OR BUPROPION OR NEFAZODONE OR MIRTAZAPINE OR VENLAFAXINE OR DULOXETINE OR MILNACIPRAN OR REBOXETINE ZIMELIDINE OR INDALPINE OR GEPIRONE OR FEMOXETINE OR ALAPROCLATE

=> s suicide or suicidal or suicidality

7625 SUICIDE
1624 SUICIDAL
191 SUICIDALITY

L4 8608 SUICIDE OR SUICIDAL OR SUICIDALITY

=> s l1 and l2 and (l3 or l4)

L5 612 L1 AND L2 AND (L3 OR L4)

=> s l5 and (PY<2003 or AY<2003 or PRY<2003)

22984586 PY<2003
4509781 AY<2003
3979358 PRY<2003

L6 148 L5 AND (PY<2003 OR AY<2003 OR PRY<2003)

=> file stnguide

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	5.70	7.46

FILE 'STNGUIDE' ENTERED AT 16:31:35 ON 25 AUG 2009
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Aug 21, 2009 (20090821/UP).

=> file hcaplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.07	7.53

FILE 'HCAPLUS' ENTERED AT 16:32:06 ON 25 AUG 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 25 Aug 2009 VOL 151 ISS 9
FILE LAST UPDATED: 24 Aug 2009 (20090824/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Jun 2009
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Jun 2009

HCAplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2009.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

The ALL, BIB, MAX, and STD display formats in the CA/CAPLUS family of databases have been updated to include new citing references information. This enhancement may impact record import into database management software. For additional information, refer to NEWS 9.

=> s depression or depressive

95095 DEPRESSION
10789 DEPRESSIVE
L7 99340 DEPRESSION OR DEPRESSIVE

=> s 16 and 17

L8 51 L6 AND L7

=> file stnguide

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	2.85	10.38

FILE 'STNGUIDE' ENTERED AT 16:32:09 ON 25 AUG 2009
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Aug 21, 2009 (20090821/UP).

=> d l8 1-51 ti
YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS' - CONTINUE? (Y)/N:y

L8 ANSWER 1 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Conjugated psychotropic drugs and uses thereof

L8 ANSWER 2 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Methods and compositions using cyclooxygenase 2 (COX-2) inhibitors for the treatment of psychiatric disorders, and combination therapies

L8 ANSWER 3 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Pharmaceutical compositions for prevention of overdose or abuse

L8 ANSWER 4 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Compositions and methods for the treatment of parkinson's disease and tardive dyskinesias with quinoline ring-containing neuromelanin-binding compounds

L8 ANSWER 5 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Carbostyryl derivatives and serotonin reuptake inhibitors for treatment of mood disorders

L8 ANSWER 6 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Combinations of medicaments comprising an alcohol deterrent for treating alcohol dependence or alcohol abuse

L8 ANSWER 7 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Method of using a cyclooxygenase 2 (COX-2) inhibitor and a 5-HT1A receptor modulator as a combination therapy for pain, inflammation, and other conditions

L8 ANSWER 8 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Stereoisomers of p-hydroxy-milnacipran, and therapeutic use

L8 ANSWER 9 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Combination therapy for depression, prevention of

suicide, and various medical and psychiatric conditions

- L8 ANSWER 10 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Combination of atypical antipsychotic and serotonin reuptake inhibitor for the treatment of chronic pain
- L8 ANSWER 11 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Single nucleotide polymorphisms (SNPs) in human DGCR2 locus and neighboring loci associated with schizophrenia and their diagnostic and therapeutic uses
- L8 ANSWER 12 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Association of SNPs in the COMT locus and neighboring loci with schizophrenia, bipolar disorder, breast cancer and colorectal cancer
- L8 ANSWER 13 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Glucocorticoid blocking agents for increasing blood-brain barrier permeability
- L8 ANSWER 14 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Methods and compositions using a cyclooxygenase 2 (COX-2) inhibitor for the treatment of psychiatric disorders
- L8 ANSWER 15 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Nefazodone in Psychotic Unipolar and Bipolar Depression : A Retrospective Chart Analysis and Open Prospective Study on Its Efficacy and Safety versus Combined Treatment with Amitriptyline and Haloperidol
- L8 ANSWER 16 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Collegium Internationale Neuro-Psychopharmacologicum (C.I.N.P.): XXIIIrd congress: Montreal, Canada, 23-27 June 2002
- L8 ANSWER 17 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Use of cyclooxygenase 2 (COX-2) inhibitors for the treatment of schizophrenia, delusional disorders, affective disorders, autism, or tic disorders
- L8 ANSWER 18 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Venlafaxine and reversible blepharoedema
- L8 ANSWER 19 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Management of symptoms associated with advanced cancer: olanzapine and mirtazapine
- L8 ANSWER 20 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Effects of psychotropic drugs on seizure threshold
- L8 ANSWER 21 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane, compositions thereof, and uses as an anti-depressant agent
- L8 ANSWER 22 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Administration of carvedilol to mitigate tardive movement disorders, psychosis, mania, and depression
- L8 ANSWER 23 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
TI Putting metabolic side effects into perspective: Risks versus benefits of atypical antipsychotics
- L8 ANSWER 24 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Severe depression: is there a best approach?
 L8 ANSWER 25 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Treatment of mood-congruent psychotic depression with imipramine
 L8 ANSWER 26 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Treatment of suicidality in schizophrenia
 L8 ANSWER 27 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Algorithm for the treatment of chronic depression
 L8 ANSWER 28 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Tablets containing 2-hydroxymethylolanzapine
 L8 ANSWER 29 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Methods and compositions for the treatment of neuroleptic and related disorders using sertindole derivatives
 L8 ANSWER 30 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Serotonergic agonists and antagonists for treatment of bronchoconstriction
 L8 ANSWER 31 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Nefazodone in the adjunctive therapy of schizophrenia: An open-label exploratory study
 L8 ANSWER 32 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI A novel approach to the identification of psychiatric drugs: serotonin-glutamate interactions in the prefrontal cortex
 L8 ANSWER 33 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Desmethylolanzapine compositions and methods
 L8 ANSWER 34 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Pharmaceutical compositions containing olanzapine-N-oxide
 L8 ANSWER 35 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI 2-Hydroxymethylolanzapine compositions and methods
 L8 ANSWER 36 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Dopamine and depression therapeutic implications
 L8 ANSWER 37 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Combination therapy of atypical antipsychotics and serotonin reuptake inhibitors for treatment of bipolar disorders
 L8 ANSWER 38 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Combination of 5-HT3 receptor antagonist and serotonin reuptake inhibitor for treatment of depression
 L8 ANSWER 39 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Atypical antipsychotic agent-serotonin reuptake inhibitor combinations for therapy of refractory depression
 L8 ANSWER 40 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Methods for treating neuropsychiatric disorders
 L8 ANSWER 41 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Efficacy of SSRIs and newer antidepressants in severe depression : comparison with TCAs

L8 ANSWER 42 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Mirtazapine: A review of its use in major depression

L8 ANSWER 43 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Pharmacotherapy for personality disorders

L8 ANSWER 44 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Olanzapine response in psychotic depression

L8 ANSWER 45 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Bupropion treatment in veterans with posttraumatic stress disorder: an open study

L8 ANSWER 46 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Atypical antipsychotics for treatment of depression in schizophrenia and affective disorders

L8 ANSWER 47 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Serotonin 5-HT₂ receptor antagonists: potential in the treatment of psychiatric disorders

L8 ANSWER 48 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Rational polypharmacy in the bipolar affective disorders

L8 ANSWER 49 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Serotonin-2 receptor-mediated intraplatelet calcium mobilization in affective disorders. Relevance to the pathophysiology of depression

L8 ANSWER 50 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Bupropion and thiothixene versus placebo and thiothixene in the treatment of depression in schizophrenia

L8 ANSWER 51 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Pharmacology in vivo of the phenylindan derivative, Lu 19005, a new potent inhibitor of dopamine, noradrenaline and 5-hydroxytryptamine uptake in rat brain

=> d 18 5 9 10 18 19 21 23 24 27 28 29 34 35 36 38 40 41 42 49 ti abs bib
 YOU HAVE REQUESTED DATA FROM FILE 'HCAPLUS' - CONTINUE? (Y)/N:y

L8 ANSWER 5 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Carbasteryl derivatives and serotonin reuptake inhibitors for treatment of mood disorders

AB The pharmaceutical composition of the present invention comprises (1) a carbasteryl derivative and (2) a serotonin reuptake inhibitor in a pharmaceutically acceptable carrier. The carbasteryl derivative may be aripiprazole or a metabolite thereof, which is a dopamine-serotonin system stabilizer. The serotonin reuptake inhibitor may be fluoxetine, duloxetine, venlafaxine, milnacipran, citalopram, fluvoxamine, paroxetine, sertraline or escitalopram. The pharmaceutical composition of the present invention is useful for treating patients with mood disorders, particularly depression or major depressive disorder. For example, a tablet formulation contained aripiprazole anhydride crystals B 5 mg, venlafaxine 75

mg, starch 131 mg, magnesium stearate 4 mg, and lactose 60 mg.
 AN 2004:589419 HCAPLUS <<LOGINID:20090825>>
 DN 141:128865
 TI Carbostyryl derivatives and serotonin reuptake inhibitors for treatment of mood disorders
 IN Kikuchi, Tetsuro; Iwamoto, Taro; Hirose, Tsuyoshi
 PA Otsuka Pharmaceutical Co., Ltd., Japan
 SO PCT Int. Appl., 92 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004060374	A1	20040722	WO 2003-JP16724	20031225 <--
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2511619	A1	20040722	CA 2003-2511619	20031225 <--
	AU 2003295235	A1	20040729	AU 2003-295235	20031225 <--
	AU 2003295235	B2	20080619		
	EP 1575590	A1	20050921	EP 2003-786308	20031225 <--
	EP 1575590	B1	20071024		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	BR 2003017771	A	20051122	BR 2003-17771	20031225 <--
	CN 1726039	A	20060125	CN 2003-80106103	20031225 <--
	EP 1723957	A2	20061122	EP 2006-17539	20031225 <--
	R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PT, RO, SE, SI, SK, TR, LT, LV				
	CN 1989968	A	20070704	CN 2007-10001620	20031225 <--
	NZ 540054	A	20070928	NZ 2003-540054	20031225 <--
	AT 376419	T	20071115	AT 2003-786308	20031225 <--
	ES 2295677	T3	20080416	ES 2003-786308	20031225 <--
	NZ 556779	A	20081224	NZ 2003-556779	20031225 <--
	RU 2356554	C2	20090527	RU 2005-123808	20031225 <--
	JP 2004217650	A	20040805	JP 2003-433429	20031226 <--
	JP 4284524	B2	20090624		
	NO 2005002359	A	20050718	NO 2005-2359	20050512 <--
	ZA 2005003873	A	20060830	ZA 2005-3873	20050513 <--
	MX 2005006857	A	20050818	MX 2005-6857	20050622 <--
	IN 2005KN01229	A	20060630	IN 2005-KN1229	20050624 <--
	KR 842694	B1	20080701	KR 2005-712073	20050624 <--
	US 20060154938	A1	20060713	US 2005-540577	20051216 <--
	KR 2007093001	A	20070914	KR 2007-717722	20070731 <--
	KR 858852	B1	20080917		
	IN 2007KN03698	A	20080125	IN 2007-KN3698	20071001 <--
PRAI	JP 2002-379003	A	20021227	<--	
	US 2003-470481P	P	20030514		
	CN 2003-80106103	A3	20031225		
	EP 2003-786308	A3	20031225		
	NZ 2003-540054	A3	20031225		
	WO 2003-JP16724	W	20031225		
	IN 2005-KN1229	A3	20050624		

KR 2005-712073 A3 20050624

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT
 OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)
 RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 9 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Combination therapy for depression, prevention of
 suicide, and various medical and psychiatric conditions
 AB The present invention relates to a new method of treatment for persons
 meeting diagnoses for major depressive disorder, or other
 unipolar (non-bipolar, nonpsychotic and non-treatment resistant)
 depression. The method comprises administering a combination of
 two categories of drugs, antipsychotics or dopamine system stabilizers, in
 combination with a newer antidepressant such as a
 selective serotonin reuptake inhibitor
 , as initial treatment or as soon as possible. The method targets the
 prevention of suicide, and provides other benefits including
 preventing disease progression development of tolerance toward the
 antidepressants. Another aspect of the invention relates to using the
 method for alleviating cognitive distortion and related functional
 impairment or health risks, and/or using the method for smoking cessation
 or nicotine withdrawal.

AN 2004:100942 HCAPLUS <<LOGINID:20090825>>

DN 140:139528

TI Combination therapy for depression, prevention of
 suicide, and various medical and psychiatric conditions

IN Migaly, Peter

PA USA

SO PCT Int. Appl., 28 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004010932	A2	20040205	WO 2003-US23326	20030725 <--
	WO 2004010932	A3	20040722		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2529857	A1	20040205	CA 2003-2529857	20030725 <--
	AU 2003268026	A1	20040216	AU 2003-268026	20030725 <--
	US 20040204401	A1	20041014	US 2003-627358	20030725 <--
	EP 1551393	A2	20050713	EP 2003-748977	20030725 <--
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	MX 2005000294	A	20050819	MX 2005-294	20050104 <--
PRAI	US 2002-319436P	P	20020730	<--	
	US 2003-627358	A	20030725		
	WO 2003-US23326	W	20030725		

OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)
 RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 10 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Combination of atypical antipsychotic and serotonin reuptake
 inhibitor for the treatment of chronic pain
 AB This invention relates to the use of the combined action of an atypical
 antipsychotic and a serotonin reuptake inhibitor for the treatment
 of chronic pain.
 AN 2003:9/1923 HCAPLUS <<LOGINID:20090825>>
 DN 140:8867
 TI Combination of atypical antipsychotic and serotonin reuptake
 inhibitor for the treatment of chronic pain
 IN Scheel-Krueger, Jorgen; Blackburn-Munro, Gordon John
 PA Neurosearch A/S, Den.
 SO PCT Int. Appl., 13 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003101492	A2	20031211	WO 2003-DK353	20030527 <--
	WO 2003101492	A3	20040129		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG			
	AU 2003227521	A1	20031219	AU 2003-227521	20030527 <--
PRAI	DK 2002-833	A	20020530	<--	
	WO 2003-DK353	W	20030527		
OSC.G	1	THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)			
RE.CNT	2	THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD			
		ALL CITATIONS AVAILABLE IN THE RE FORMAT			

L8 ANSWER 18 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Venlafaxine and reversible blepharoadema
 AB The newer antidepressant venlafaxine is known to cause
 dilutional hyponatremia, but to our knowledge no reports on localized
 edemas in the absence of electrolyte disturbances are available. We
 present a case in which venlafaxine caused reversible
 blepharoadema in an otherwise phys. healthy patient. Ms. M., a 25-yr-old
 woman, suffered from schizoaffective disorder since being 23 yr old. Upon
 administration of quetiapine, she completely recovered but
 relapsed twice due to medical non-compliance, resulting in the third
 hospitalization. Again, psychotic symptoms cleared upon prescription of
 600 mg quetiapine; further, 45 mg mirtazapine was
 given. Quetiapine remained at a stable dose for 10 wk,
 mirtazapine for 2 wk; no side-effects were reported by the patient
 or observed by her physicians and no other medication was used. As she
 persistently complained about depressed mood, loss of motivation and
 drive, we adminl. administered 75 mg of retarded venlafaxine in
 the morning. The next day, marked bilateral and sym. blepharoadema could
 be noted which did not ache on palpation, but caused discomfort on eye
 movements, generally worrying Ms. M. She had no relevant past medical
 history besides her psychiatric disorder, especially no occurrence of allergic
 sensitivity, and had never experienced localized edema. No other edemas

were present, nor were other medical symptoms. Serum electrolytes were within the normal range. Believing that venlafaxine caused lid edema, we discontinued venlafaxine after the second day; within 24 h, the symptom completely vanished.

AN 2002:929206 HCAPLUS <<LOGINID::20090825>>

DN 140:139166

TI Venlafaxine and reversible blepharoidema

AU Reif, Andreas; Pfuhlmann, Bruno

CS Department of Psychiatry, Julius-Maximilians-University of Wuerzburg, Wuerzburg, 97080, Germany

SO International Journal of Neuropsychopharmacology (2002), 5(4), 413-414

CODEN: IJNUFB; ISSN: 1461-1457

PB Cambridge University Press

DT Journal

LA English

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 19 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Management of symptoms associated with advanced cancer: olanzapine and mirtazapine

AB A review. Advanced cancer patients are polysymptomatic and often receive multiple medications for symptom relief. Common symptoms include anorexia, weight loss, delirium and depression. Olanzapine and mirtazapine may have several advantages over older agents despite increased acquisition costs. Both medications can treat several symptoms with a low risk for drug-drug interactions and with only once- or twice-daily dosing. Drug side effects are low, compared with more conventionally used agents. The pharmacokinetics and pharmacodynamics of both agents are unique and explain many of the benefits. More research and clin. experience will be necessary to define their role in the palliation of advanced cancer.

AN 2002:720957 HCAPLUS <<LOGINID::20090825>>

DN 137:272678

TI Management of symptoms associated with advanced cancer: olanzapine and mirtazapine

AU Davis, Mellar P.; Khawam, Elias; Pozuelo, Leo; Lagman, Ruth

CS Harry R. Horvitz Cent. for Palliative Med., Cleveland Clin. Found., Cleveland, OH, 44195, USA

SO Expert Review of Anticancer Therapy (2002), 2(4), 365-376

CODEN: ERATBJ; ISSN: 1473-7140

PB Future Drugs Ltd.

DT Journal; General Review

LA English

OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

RE.CNT 88 THERE ARE 88 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 21 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane, compositions thereof, and uses as an anti-depressant agent

AB The present invention relates to (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane and pharmaceutically acceptable salts thereof, comps. comprising (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane or a pharmaceutically acceptable salt thereof, and methods for treating or preventing depression in a patient comprising administering (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane or a pharmaceutically acceptable salt thereof. The (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane or pharmaceutically acceptable salt thereof is preferably substantially free of its corresponding (-)-enantiomer. The +

isomer is obtained by HPLC resolution on a CHIRALPAK AD column. The + isomer has greater affinity for both norepinephrine and serotonin uptake sites in rat forebrain membranes than the ± compound. The + isomer is administered along with a known antidepressant, anxiolytic, antipsychotic or antiobesity agent in treatment of various depression conditions including depression associated with anxiety, seizures, menopause, alcoholism, etc.

AN 2002:290820 HCAPLUS <<LOGINID:20090825>>

DN 136:304102

TI (+)-1-(3,4-dichlorophenyl)-3-azabicyclo[3.1.0]hexane, compositions thereof, and uses as an anti-depressant agent

IN Lipa, Arnold Stan; Epstein, Joseph William

PA Dov Pharmaceutical, Inc., USA

SO U.S., 7 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 4

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6372919	B1	20020416	US 2001-758883	20010111 <--
	CA 2434616	A1	20020829	CA 2002-2434616	20020111 <--
	WO 2002066427	A2	20020829	WO 2002-US845	20020111 <--
	WO 2002066427	A3	20030313		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2002251758	A1	20020904	AU 2002-251758	20020111 <--
	AU 2002251758	B2	20080103		
	EP 1349835	A2	20031008	EP 2002-720783	20020111 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	HU 2003002613	A2	20031128	HU 2003-2613	20020111 <--
	HU 2003002613	A3	20070928		
	BR 2002006434	A	20031230	BR 2002-6434	20020111 <--
	CN 1496349	A	20040512	CN 2002-806351	20020111 <--
	ZA 2003005440	A	20040715	ZA 2003-5440	20020111 <--
	JP 2005500983	T	20050113	JP 2002-565944	20020111 <--
	NZ 527101	A	20050826	NZ 2002-527101	20020111 <--
	RU 2294926	C2	20070310	RU 2003-124649	20020111 <--
	CN 101461804	A	20090624	CN 2008-10185945	20020111 <--
	NO 2003003165	A	20030904	NO 2003-3165	20030710 <--
	NO 325709	B1	20080707		
	MX 2003006210	A	20041015	MX 2003-6210	20030711 <--
	IN 2003CN01224	A	20051118	IN 2003-CN1224	20030807 <--
	US 20040132797	A1	20040708	US 2004-466457	20040210 <--
	US 7098229	B2	20060829		
PRAI	US 2001-758883	A	20010111	<--	
	CN 2002-806351	A3	20020111	<--	
	WO 2002-US845	W	20020111	<--	

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 23 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Putting metabolic side effects into perspective: Risks versus benefits of
 atypical antipsychotics
 AB A review. The lengthy list of the side effects and morbidity associated with
 the atypical antipsychotics might make a patient with psychosis and his or
 her caregivers so concerned about the use of any of these medications,
 particularly those associated with a higher risk of diabetes, weight gain, or
 increased lipid levels, that they would prefer to avoid all of them.
 However, schizophrenia is associated with a relatively high risk for several
 diseases, including diabetes, that is independent of the risks that are
 linked to atypical antipsychotic use. Therefore, the clinician
 who might think, "Why use atypicals if using the typical drugs will escape
 the problems of monitoring and all the associated effects of diabetes and
 hyperglycemia" needs to know that these problems cannot be avoided simply
 by choosing typical antipsychotics. Clinicians, patients, and concerned
 family members must balance the significant benefits of atypical
 antipsychotic treatment - improved cognition, reduced
 suicidality, and less depression - against the risks of
 metabolic disturbances and select a course of treatment that includes a
 realistic monitoring program.
 AN 2002:75124 HCAPLUS <<LOGINID::20090825>>
 DN 136:272542
 TI Putting metabolic side effects into perspective: Risks versus benefits of
 atypical antipsychotics
 AU Meltzer, Herbert Y.
 CS Department of Psychiatry and Pharmacology, Division of Psychopharmacology,
 Vanderbilt University Medical Center, Nashville, TN, 37212, USA
 SO Journal of Clinical Psychiatry (2001), 62(Suppl. 27), 35-39
 CODEN: JCLPDE; ISSN: 0160-6689
 PB Physicians Postgraduate Press, Inc.
 DT Journal; General Review
 LA English
 OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)
 RE.CNT 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 24 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Severe depression: is there a best approach?
 AB A review. A major depressive episode can be categorized as
 severe based on depressive symptoms, scores on
 depression rating scales, the need for hospitalization,
 depressive subtypes, functional capacity, level of
 suicidality and the impact that the depression has on
 the patient. Several biol., psychol. and social factors, and the presence
 of comorbid psychiatric or medical illnesses, impact on depression
 severity. A number of factors are reported to influence outcome in severe
 depression, including duration of illness before treatment,
 severity of the index episode, treatment modality used, and dosage and
 duration of and compliance with treatment. Potential complications of
 untreated severe depression include suicide,
 self-mutilation and refusal to eat, and treatment resistance. Several
 antidepressants have been studied in the treatment of severe
 depression. These include tricyclic antidepressants (TCAs),
 selective serotonin reuptake inhibitors (SSRIs), serotonin-noradrenaline
 (norepinephrine) reuptake inhibitors, noradrenergic and specific
 serotonergic antidepressants, serotonin 5-HT2 receptor antagonists,
 monoamine oxidase inhibitors, and amfebutamone (bupropion).
 More recently, atypical antipsychotics have shown some utility in the
 management of severe and resistant depression. Data on the
 differential efficacy of TCAs vs. SSRIs and the newer antidepressants in
 severe depression are mixed. Some studies have reported that

TCAs are more efficacious than SSRIs; however, more recent studies have shown that TCAs and SSRIs have equivalent efficacy. There are reports that some of the newer antidepressants may be more effective than SSRIs in the treatment of severe depression, although the sample sizes in some of these studies were small. Combination therapy has been reported to be effective. The use of an SSRI-TCA combination, while somewhat controversial, may rapidly reduce depressive symptoms in some patients with severe depression. The combination of an antidepressant and an antipsychotic drug is promising and may be considered for severe depression with psychotic features. Although the role of cognitive behavior therapy (CBT) in severe depression has not been adequately studied, a trial of CBT may be considered in severely depressed patients whose symptoms respond poorly to an adequate antidepressant trial, who are intolerant of antidepressants, have contraindications to pharmacotherapy, and who refuse medication or other somatic therapy. A combination of CBT and antidepressants may also be beneficial in some patients. Electroconvulsive therapy (ECT) may be indicated in severe psychotic depression, severe melancholic depression, resistant depression, and in patients intolerant of antidepressant medications and those with medical illnesses which contraindicate the use of antidepressants (e.g. renal, cardiac or hepatic disease).

AN 2001:908128 HCAPLUS <<LOGINID::20090825>>

DN 136:193477

TI Severe depression: is there a best approach?

AU Sonawalla, Shamsah B.; Fava, Maurizio

CS Depression Clinical and Research Program, Harvard Medical School, Massachusetts General Hospital, Boston, MA, USA

SO CNS Drugs (2001), 15(10), 765-776

CODEN: CNDREF; ISSN: 1172-7047

PB Adis International Ltd.

DT Journal; General Review

LA English

OSC.G 21 THERE ARE 21 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)

RE.CNT 105 THERE ARE 105 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 27 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Algorithm for the treatment of chronic depression

AB A review with 41 refs. Chronic depression, which is marked by a course of illness lasting 2 yr or more, encompasses 4 subtypes of depressive illness: (1) chronic major depressive disorder, (2) dysthymic disorder, (3) dysthymic disorder with major depressive disorder ("double depression"), and (4) major depressive disorder with poor interepisodic recovery (i.e., in incomplete remission). In the 1990s, chronic depression had a reported prevalence rate of 3% to 5% and accounted for 30% to 35% of all cases of depression in the United States. The authors present an algorithm modified from the Texas Medication Algorithm Project for patients with chronic depression. This treatment algorithm recommends a progression of steps or stages in treating chronic depression. The first stage is monotherapy with the selective serotonin reuptake inhibitors, nefazodone, bupropion sustained release, venlafaxine extended release, mirtazapine, or psychotherapy. Later options include combination therapy, electroconvulsive therapy, atypical antipsychotics, and novel treatments. Utilization of a comprehensive treatment algorithm for chronic major depression should encourage efficient, efficacious treatment.

AN 2001:311359 HCAPLUS <<LOGINID::20090825>>

DN 135:220442

TI Algorithm for the treatment of chronic depression
 AU Trivedi, Madhukar H.; Kleiber, Beverly A.
 CS Depression and Anxiety Disorders Program, Southwestern Medical Center at
 Dallas, The University of Texas, Dallas, TX, 75390-9101, USA
 SO Journal of Clinical Psychiatry (2001), 62(Suppl. 6), 22-29
 CODEN: JCLPDE; ISSN: 0160-6689
 PB Physicians Postgraduate Press, Inc.
 DT Journal; General Review
 LA English
 OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)
 RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 28 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Tablets containing 2-hydroxymethylolanzapine
 AB Methods and compns. are disclosed utilizing 2-hydroxymethylolanzapine (I)
 for the treatment of psychosis in humans. I exhibits a low tendency
 toward drug-drug interactions and a more predictable dosing regimen than
 olanzapine. I is also useful for the treatment of acute mania,
 mild anxiety states, anxiety disorders, schizophrenia, bipolar disorder,
 attention deficit hyperactivity disorder, autistic disorder, excessive
 aggression, substance abuse, depressive signs and symptoms, tic
 disorder, functional bowel disorder and fungal dermatitis. Thus, tablets
 contained I 20, croscarmellose 60, colloidal SiO2 8, Mg stearate 1,
 microcryst. cellulose 190, Croscarmellose 15, and talc 10 mg.

AN 2001:45171 HCAPLUS <<LOGINID::20090825>>
 DN 134:91165
 TI Tablets containing 2-hydroxymethylolanzapine
 IN Yelle, William E.
 PA Sepracor Inc., USA
 SO U.S., 6 pp.
 CODEN: USXXAM
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 6174882	B1	20010116	US 1999-444160	19991122 <--
	US 6346528	B1	20020212	US 2000-690357	20001017 <--
PRAI	US 1998-109552P	P	19981123	<--	
	US 1999-444160	A3	19991122	<--	
OSC.G	1				THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)
RE.CNT	36				THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 29 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Methods and compositions for the treatment of neuroleptic and related
 disorders using sertindole derivatives
 AB The invention relates to novel methods using, and pharmaceutical compns.
 and dosage forms comprising, sertindole derivs. Sertindole derivs.
 include, but are not limited to, nor-sertindole, 5-oxo-sertindole,
 dehydro-sertindole, and dehydro-nor-sertindole. The methods of the
 invention are directed to the treatment and prevention of neuroleptic and
 related disorders such as, but are not limited to, psychotic disorders,
 depression, anxiety, substance addiction, memory impairment and
 pain. For example, capsules were prepared containing a sertindole derivative
 50.0 mg, lactose 48.5 mg, TiO2 0.5 mg, and Mg stearate 1.0 mg.
 AN 2000:861482 HCAPLUS <<LOGINID::20090825>>
 DN 134:32977
 TI Methods and compositions for the treatment of neuroleptic and related

disorders using sertindole derivatives
 IN Jerussi, Thomas P.
 PA Sepracor Inc., USA
 SO PCT Int. Appl., 33 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000072837	A2	20001207	WO 2000-US14984	20000531 <--
	WO 2000072837	A3	20010517		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, NI, TD, TG			
	US 6489341	B1	20021203	US 2000-580492	20000530 <--
PRAI	US 1999-137447P	P	19990602	<--	
	US 2000-580492	A	20000530	<--	

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT
 OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)
 RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 34 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Pharmaceutical compositions containing olanzapine-N-oxide
 AB Methods and compns. are disclosed utilizing olanzapine-N-oxide for the treatment of psychosis in humans. Olanzapine-N-oxide exhibits a lessened liability toward drug-drug interactions than olanzapine and a more predictable dosing regimen than olanzapine. Olanzapine-N-oxide is also useful for the treatment of acute mania, mild anxiety states, anxiety disorders, schizophrenia, bipolar disorder, attention deficit hyperactivity disorder, autistic disorder, excessive aggression, substance abuse, depressive signs and symptoms, tic disorder, functional bowel disorder and fungal dermatitis. The invention also relates to pharmaceutical compns. comprising olanzapine-N-oxide. E.g., preparation of tablets containing olanzapine-N-oxide 10 and 20 mg was described.

AN 2000:577484 HCAPLUS <<LOGINID:20090825>>
 DN 133:144934
 TI Pharmaceutical compositions containing olanzapine-N-oxide
 IN Yelle, William E.
 PA Sepracor Inc., USA
 SO PCT Int. Appl., 21 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000030649	A1	20000602	WO 1999-US27644	19991122 <--
	W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,			

SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE,
 DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,
 CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

CA 2351719 A1 20000602 CA 1999-2351719 19991122 <--
 US 6121259 A 20000919 US 1999-444159 19991122 <--
 EP 1135136 A1 20010926 EP 1999-961750 19991122 <--
 EP 1135136 B1 20031105

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO

JP 2002530340 T 20020917 JP 2000-583532 19991122 <--
 AU 757870 B2 20030306 AU 2000-18266 19991122 <--
 AT 253364 T 20031115 AT 1999-961750 19991122 <--
 ES 2211205 T3 20040701 ES 1999-961750 19991122 <--
 US 6352984 B1 20020305 US 2000-632584 20000807 <--
 US 20020065272 A1 20020530 US 2001-16205 20011030 <--

PRAI US 1998-109551P F 19981123 <--
 US 1999-444159 A3 19991122 <--
 WO 1999-US27644 W 19991122 <--
 US 2000-632584 A3 20000807 <--

OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)
 RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 35 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI 2-Hydroxymethylolanzapine compositions and methods
 AB Methods and comps. are disclosed using 2-hydroxymethylolanzapine for the
 treatment of psychosis in humans. 2-Hydroxymethylolanzapine exhibits a
 lessened liability toward drug-drug interactions than olanzapine
 and a more predictable dosing regimen than olanzapine.
 2-Hydroxymethylolanzapine is also useful for the treatment of acute mania,
 mild anxiety states, anxiety disorders, schizophrenia, bipolar disorder,
 attention deficit hyperactivity disorder, autistic disorder, excessive
 aggression, substance abuse, depressive signs and symptoms, tic
 disorder, functional bowel disorder and fungal dermatitis.

AN 2000:577483 HCAPLUS <<LOGINID:20090825>>
 DN 133:144933
 TI 2-Hydroxymethylolanzapine compositions and methods
 IN Yelle, William E.
 PA Sepracor Inc., USA
 SO PCT Int. Appl., 22 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000030648	A1	20000602	WO 1999-US27640	19991122 <--
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	CA 2352611 A1 20000602			CA 1999-2352611	19991122 <--
	EP 1133299 A1 20010919			EP 1999-959066	19991122 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	JP 2002530339 T 20020917			JP 2000-583531	19991122 <--

AU 757874 B2 20030306 AU 2000-16315 19991122 <--
 PRAI US 1998-109552P P 19981123 <--
 WO 1999-US27640 W 19991122 <--
 OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)
 RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 36 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Dopamine and depression therapeutic implications
 AB A review with 83 refs. The mesolimbic dopaminergic system functions as a major reward pathway in the CNS and is an appropriate target for antidepressant drugs. This review describes the principal features of drugs such as amfebutamone (bupropion) which activate the mesolimbic system without inducing strong neuroadaptation and are therefore useful in the treatment of retarded (or inhibited) depression. The short latency of clin. action makes these drugs particularly suitable for the treatment of patients with severe depression or those who are poorly compliant with other medications. Addnl. dopaminergic antidepressants include minaprine and amisulpride. The latter drug potentiates dopaminergic transmission through an atypical mechanism, i.e., the inhibition of dopamine autoreceptors controlling the synthesis and release of dopamine. Finally, a number of drugs that are not considered as classical "dopaminergic" antidepressants, such as tricyclic antidepressants or selective serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitors, can also affect dopaminergic transmission, as indicated for example by their ability to induce changes in brain dopamine receptor d. This further supports the importance of central dopaminergic transmission in the pathophysiol. of depression.

2000:92738 HCAPLUS <<LOGINID::20090825>>
 DN 132:232035
 TI Dopamine and depression therapeutic implications
 AU Rampello, Liborio; Nicoletti, Ferdinando; Nicoletti, Francesco
 CS Institute of Neurological Sciences, Policlinico Universitario, University of Catania, Catania, Italy
 SO CNS Drugs (2000), 13(1), 35-45
 CODEN: CNDREF; ISSN: 1172-7047
 PB Adis International Ltd.
 DT Journal; General Review
 LA English
 OSC.G 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)
 RE.CNT 83 THERE ARE 83 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 38 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Combination of 5-HT3 receptor antagonist and serotonin reuptake inhibitor for treatment of depression
 AB The present invention provides a method for treating a patient suffering from depression, comprising administering to said patient an effective amount of a first component which is a 5-HT3 receptor antagonist, in combination with an effective amount of a second component which is a serotonin reuptake inhibitor wherein improvement in sexual dysfunction and/or reduction in gastrointestinal side effects is recognized. Various formulations were prepared. E.g., a tablet was prepared using zatosetron 10, fluoxetine HCl 10, microcryst. cellulose 275, fumed silica 10, and stearic acid 5 mg, resp.
 AN 1999:753081 HCAPLUS <<LOGINID::20090825>>
 DN 131:346552
 TI Combination of 5-HT3 receptor antagonist and serotonin reuptake inhibitor for treatment of depression
 IN Michelson, David; Tollefson, Gary Dennis

PA Eli Lilly and Company, USA
 SO PCT Int. Appl., 25 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 9959593	A1	19991125	WO 1999-US10092	19990510 <--
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2332253	A1	19991125	CA 1999-2332253	19990510 <--
AU 9938912	A	19991206	AU 1999-38912	19990510 <--
EP 1077704	A1	20010228	EP 1999-921795	19990510 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT, LV, FI, RO			
JP 2002515435	T	20020528	JP 2000-549258	19990510 <--
PRAI US 1998-86268P	P	19980521	<--	
WO 1999-US10092	W	19990510	<--	
OSC.G 6	THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)			
RE.CNT 7	THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD			
	ALL CITATIONS AVAILABLE IN THE RE FORMAT			

L8 ANSWER 40 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Methods for treating neuropsychiatric disorders

AB The invention provides methods for treating neuropsychiatric disorders such as schizophrenia, Alzheimer's Disease, autism, depression, benign forgetfulness, childhood learning disorders, cluse head injury, and attention deficit disorder. The methods entail administering to a patient with a neuropsychiatric disorder a pharmaceutical composition containing (i) a therapeutically effective amount of D-alanine (or a modified form), provided that the composition is substantially free of D-cycloserine, and/or (ii) D-serine (or a modified form), and/or (iii) 105 to 500 mg of D-cycloserine (or a modified form), and/or (iv) N-methylglycine (or a modified form). Using double-blind conditions, patients were randomly assigned to receive placebo (fruit juice), D-serine 30, D-alanine 60-100, or N-methylglycine 30 mg/kg/day once a day by mouth for 6 wk. Treatment with D-serine, D-alanine, or N-methylglycine improved the schizophrenic symptoms and cognitive deficit of the patients. Specifically, treatment with D-serine resulted in a 21% reduction of the neg. symptoms (on the SANS scale), and it resulted in a 17% reduction of the pos. symptoms. Treatment with D-alanine resulted in an 11% reduction of the neg. symptoms and a 12% reduction of the

pos. symptoms. Reatment with N-methylglycine resulted in a 20% reduction of the neg. symptoms and a 15% reduction of the pos. symptoms. These redns. in the neg. and pos. symptoms represented clin. significant improvement.

AN 1999:672562 HCAPLUS <<LOGINID:20090825>>

DN 131:281590

TI Methods for treating neuropsychiatric disorders

IN Tsai, Guochuan; Coyle, Joseph

PA The General Hospital Corporation, USA

SO PCT Int. Appl., 27 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9952519	A2	19991021	WO 1999-US8056	19990414 <--
	WO 9952519	A3	19991202		
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	CA 2328197	A1	19991021	CA 1999-2328197	19990414 <--
	CA 2328197	C	20071120		
	CA 2601132	A1	19991021	CA 1999-2601132	19990414 <--
	AU 9935571	A	19991101	AU 1999-35571	19990414 <--
	AU 765603	B2	20030925		
	EP 1073432	A2	20010207	EP 1999-917453	19990414 <--
	EP 1073432	B1	20070815		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
	US 6228875	B1	20010508	US 1999-291296	19990414 <--
	HU 2001001627	A2	20011028	HU 2001-1627	19990414 <--
	HU 2001001627	A3	20030228		
	JP 2002511409	T	20020416	JP 2000-543129	19990414 <--
	RU 2219924	C2	20031227	RU 2000-128654	19990414 <--
	NZ 508160	A	20040130	NZ 1999-508160	19990414 <--
	IL 139008	A	20060221	IL 1999-139008	19990414 <--
	AT 369848	T	20070915	AT 1999-917453	19990414 <--
	EP 1844769	A2	20071017	EP 2007-75595	19990414 <--
	R: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
	ES 2164040	T3	20080201	ES 1999-917453	19990414 <--
	MX 2000010009	A	20010521	MX 2000-10009	20001013 <--
	US 20020035145	A1	20020321	US 2001-834351	20010413 <--
	US 6420351	B2	20020716		
	HK 1036583	A1	20080606	HK 2001-105482	20010807 <--
	US 20020193429	A1	20021219	US 2002-196686	20020715 <--
	US 6667297	B2	20031223		
	US 20040092530	A1	20040513	US 2003-668583	20030923 <--
	US 6974821	B2	20051213		
	US 20050250851	A1	20051110	US 2005-175832	20050705 <--
PRAI	US 1998-81645P	P	19980414	<--	
	US 1998-81654P	P	19980414	<--	
	CA 1999-2328197	A3	19990414	<--	
	EP 1999-917453	A3	19990414	<--	
	US 1999-291296	A1	19990414	<--	
	WO 1999-US8056	W	19990414	<--	
	US 2001-834351	A1	20010413	<--	
	US 2002-196686	A1	20020715	<--	
	US 2003-668583	A1	20030923		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OSC.G 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (21 CITINGS)

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 41 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Efficacy of SSRIs and newer antidepressants in severe depression : comparison with TCAs

AB A review with 58 refs. The significant morbidity and mortality associated

with severe depression and its psychotic or melancholic subtypes necessitate effective and well-tolerated therapy. This review evaluates antidepressant treatments for patients with severe depression. Comparative clin. trials conducted on patients with severe depression were found by an English-language MEDLINE search (1985 to present). Addnl. studies were identified in article bibliogs. Search terms included depressive disorders, depression and severe, hospitalized, melancholic or melancholia, psychotic, and endogenous. Evidence for efficacy of SSRIs in severe or melancholic depression comes from a small but growing number of controlled studies with adequate samples, as well as meta-analyses and retrospective subgroup anal. of premarketing trials. In studies that defined response as a 50% or greater reduction in Hamilton Rating Scale for Depression (HAM-D) scores, response rates ranged from 53% to 64% for SSRIs and 43% to 70% for TCAs. In sep. trials on severe depression, venlafaxine and mirtazapine were both more effective than placebo and an active comparator. Nefazodone and bupropion were each found to be more effective than placebo in studies of severe depression. Venlafaxine and mirtazapine have been found to be more effective than fluoxetine. SSRIs and TCAs are comparably effective for the treatment of severe or melancholic depression. SSRIs and other newer agents appear to be better tolerated than TCAs, specifically lacking adverse anticholinergic and cardiovascular effects that may limit the use of TCAs. Emerging data with venlafaxine and mirtazapine in severely depressed patients with or without melancholia support the efficacy of these treatments. Nefazodone and bupropion were found to be effective in hospitalized depressed patients. Electroconvulsive therapy (ECT) or combined antidepressant therapy may be useful in some patients with severe depression. Patients with severe psychotic depression may respond better to an antipsychotic-antidepressant combination.

AN 1999:402615 HCAPLUS <<LOGINID:20090825>>
 DN 131:82427
 TI Efficacy of SSRIs and newer antidepressants in severe depression
 : comparison with TCAs
 AU Hirschfeld, Robert M. A.
 CS Department of Psychiatry and Behavioral Sciences, The University of
 Texas-Medical Branch, Galveston, TX, 77555, USA
 SO Journal of Clinical Psychiatry (1999), 60(5), 326-335
 CODEN: JCLPDE; ISSN: 0160-6689
 PB Physicians Postgraduate Press, Inc.
 DT Journal; General Review
 LA English
 OSC.G 28 THERE ARE 28 CAPLUS RECORDS THAT CITE THIS RECORD (28 CITINGS)
 RE.CNT 58 THERE ARE 58 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 42 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN
 TI Mirtazapine: A review of its use in major depression
 AB A review with 107 refs. Mirtazapine is a noradrenergic and
 specific serotonergic antidepressant which has been evaluated
 predominantly in the treatment of major depression. The drug
 had efficacy equivalent to that of tricyclic antidepressants and it was at
 least as effective as trazodone in the majority of available short-term
 trials in patients with moderate or severe depression, including
 those with basal anxiety symptoms or sleep disturbance and the elderly. A
 continuation study also showed that sustained remission rates were higher
 with mirtazapine than with amitriptyline and that the drugs had
 similar efficacy for the prevention of relapse. There is some evidence

for a faster onset of action with mirtazapine than with the selective serotonin reuptake inhibitors (SSRIs). Mirtazapine was more effective than the SSRI fluoxetine after 3 and 4 wk of therapy and it was also more effective than paroxetine and citalopram after 1 and 2 wk, resp., in short-term assessments (6 or 8 wk). Preliminary data suggest that the drug may be effective as an augmentation or combination therapy in patients with refractory depression. Anticholinergic events and other events, including tremor and dyspepsia, are less common with mirtazapine than with tricyclic antidepressants. There was a greater tendency for SSRI-related adverse events with fluoxetine than with mirtazapine, but, overall, mirtazapine had a tolerability profile similar to that of the SSRIs. Increased appetite and body-weight gain appear to be the only events that are reported more often with mirtazapine than with comparator antidepressants. In vitro and in vivo data have suggested that mirtazapine is unlikely to affect the metabolism of drugs metabolized by cytochrome P 450 (CYP) 2D6, although few formal drug-interaction data are available. Conclusions: Mirtazapine is effective and well tolerated for the treatment of patients with moderate to severe major depression. Further research is required to define the comparative efficacy of mirtazapine in specific patient groups, including the elderly and those with severe depression. Clarification of its efficacy as an augmentation therapy and in patients with refractory depression and its role in improving the efficacy and reducing the extrapyramidal effects of antipsychotic drugs would also help to establish its clin. value. The low potential for interaction with drugs that are metabolized by CYP2D6, including antipsychotics, tricyclic antidepressants and some SSRIs, may also make mirtazapine an important option for the treatment of major depression in patients who require polytherapy. Mirtazapine also appears to be useful in patients with depression who have anxiety symptoms and sleep disturbance.

AN 1999:307238 HCAPLUS <<LOGINID:20090825>>
 DN 130:332190
 TI Mirtazapine: A review of its use in major depression
 AU Holm, Kristin J.; Markham, Anthony
 CS Adis International Limited, Auckland, N. Z.
 SO Drugs (1999), 57(4), 607-631

CODEN: DRUGAY; ISSN: 0012-6667

PB Adis International Ltd.

DT Journal; General Review

LA English

OSC.G 51 THERE ARE 51 CAPLUS RECORDS THAT CITE THIS RECORD (51 CITINGS)

RE.CNT 107 THERE ARE 107 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 49 OF 51 HCAPLUS COPYRIGHT 2009 ACS on STN

TI Serotonin-2 receptor-mediated intraplatelet calcium mobilization in affective disorders. Relevance to the pathophysiology of depression

AB 5-HT-stimulated intracellular Ca concentration change was studied in the platelets of healthy subjects, using fluorescent Ca indicator fura-2. 5-HT increased the Ca response in a concentration-dependent manner. The

maximal

response was obtained at 10 μ M of 5-HT and its EC50 value was 0.4 μ M. This response was potentially inhibited by selective 5-HT2 receptor antagonists, suggesting that the 5-HT-induced Ca mobilization is mediated by 5-HT2 receptors. This 5-HT-stimulated Ca response was not significantly affected by the time of blood sampling, gender, age, meal, or exercise. Therefore, it may be concluded that the 5-HT-induced Ca response in human platelets is a stable parameter and that it is suitable

for assessing 5-HT₂ receptor function in depressed patients. Thus, the 5-HT-induced Ca mobilization was measured in the platelets of depressed patients. The response was significantly higher in unmedicated patients with bipolar depression and melancholic major depression than in those with nonmelancholic major depression and normal controls. The enhanced Ca response to 5-HT failed to correlate with severity of depressive symptoms. In patients with bipolar depression and melancholic major depression, there was no significant difference in 5-HT-stimulated Ca response between unmedicated group and euthymic-treated group. These results suggest that 5-HT₂ receptor function is increased in some type of affective disorders and that the enhanced Ca response to 5-HT may be trait dependent rather than state dependent.

AN 1993:469623 HCAPLUS <<LOGINID::20090825>>

DN 119:69623

OREF 119:12541a,12544a

TI Serotonin-2 receptor-mediated intraplatelet calcium mobilization in affective disorders. Relevance to the pathophysiology of depression

AU Kusumi, Ichiro

CS Sch. Med., Hokkaido Univ., Sapporo, 060, Japan

SO Hokkaido Igaku Zasshi (1993), 68(3), 325-36

CODEN: HOIZAK; ISSN: 0367-6102

DT Journal

LA Japanese

OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)